

- Sato, T. (2002) Description of plesiosaurs (Reptilia: Sauropterygia) from the Bearpaw Formation (Campanian–Maastrichtian) and a phylogenetic analysis of the Elasmosauridae. Unpublished D. Phil. Thesis, University of Calgary.
- Welles, S. P. (1943) Elasmosaurid plesiosaurs with a description of new material from California and Colorado. University of California Publications in Geological Sciences 13: 125–215.
- Welles, S. P. (1970) The longest neck in the ocean. University of Nebraska News, Vol. 50, No. 9, 2 p.
-

**A REASSESSMENT OF HISTORICAL PLESIOSAURIAN SPECIMENS
FROM THE TURONIAN (LOWER UPPER CRETACEOUS) OF THE OPOLE AREA,
SOUTHWEST POLAND**

Sven **Sachs**^{1,2}, Robert **Niedźwiedzki**³, Mariusz **Kędzierski**⁴, Benjamin P. **Kear**⁵, Elena **Jagt-Yazykova**⁶ & John W.M. **Jagt**⁷

¹Naturkundemuseum Bielefeld, Abteilung Geowissenschaften, Adenauerplatz 2, 33602 Bielefeld, Germany [Sachs.Pal@gmail.com]

²Im Hof 9, 51766 Engelskirchen, Germany

³Institute of Geological Sciences, Wrocław University; pl. M. Bornha 9, 50-204 Wrocław, Poland [robert.niedzwiedzki@uwr.edu.pl]

⁴Institute of Geological Sciences, Jagiellonian University, Oleandry 2a, 30-063 Kraków, Poland [mariusz.kedzierski@uj.edu.pl]

⁵Museum of Evolution, Uppsala University, Norbyvägen 18, 752 36 Uppsala, Sweden [benjamin.kear@geo.uu.se]

⁶Laboratory of Palaeobiology, University of Opole, Oleska 22, 45-052 Opole, Poland

[elenajagt@gmail.com]

⁷Natuurhistorisch Museum Maastricht, de Bosquetplein 6-7, 6211 KJ Maastricht, the Netherlands [John.Jagt@maastricht.nl]

In 1897, Richard Leonhard presented brief descriptions and some illustrations of a few plesiosaurian remains (teeth and a supposed phalanx) from the lower Turonian of the Opole area (southwest Poland). Leonhard assigned the teeth to *Polyptychodon interruptus* and established a new genus, *Plesiosauridarum* (a species name was not supplied, making this a *nomen nudum*), for the supposed phalanx. Until recently, the whereabouts of all these specimens were unknown. However, during a recent survey of several Polish and German collections, most of Leonhard's specimens could be relocated. Furthermore, other historical, still undescribed specimens (i.e., a tooth and paddle element) were encountered, and these supplement Leonhard's material. Most of the original material has probably been found in now abandoned quarries in the area. Active quarrying is done at the Odra Nowa quarry in the Opole city area, but so far no plesiosaurian remains have been collected there. In total, four slender pliosauiromorph teeth from the Opole area are available for study. These differ from the typically more massive teeth of Late Cretaceous pliosaurids (*Brachauchenius*/*Megacephalosaurus*) and the pliosauiromorph teeth from coeval strata in the Bohemian Cretaceous Basin (Czech Republic; see Kear *et al.*, 2014). However, a closely similar tooth crown has recently been described from the upper Turonian of the Saxonian Cretaceous Basin (eastern Germany; see Sachs *et al.*, 2016). Generic placement of isolated pliosauiromorph teeth from Cretaceous strata to the genus *Polyptychodon* is commonly seen in old literature sources. However, this taxon is, in fact, a *nomen dubium* (see Madzia, 2015). *Polyptychodon* has long been considered to be a late pliosaurid, but the slenderness of some referred tooth crowns, including those from Opole, indicate that these might rather derive from polycotylids (see Sachs *et al.*, 2016). The presence of similar teeth in Turonian strata in Germany and Poland indicate that some pliosauiromorph taxa had a wider distribution than previously recognised.

The paddle element (probably a mesopodial) is not diagnostic at the family level and Leonhard's *Plesiosauridarum* (which, so far, could not be traced) is a mosasauroid vertebra. In order to date these early finds, powdered rock samples for calcareous nannoplankton studies were scraped from the matrix or the infill of dentin canals. Light microscope studies have demonstrated nannofossil assemblages of 32 taxa, dominated by *Watznaueria barnesae*. Based on the presence of the stratigraphically youngest species, *Quadrum gartneri*, the samples appear to be not younger than nannofossil zone UC7, i.e., probably represent the lower (lower middle?) Turonian (see Burnett, 1998).

References

- Burnett, J.A. (1998) Upper Cretaceous. In: Bown, P.R. (ed.), *Calcareous Nannofossil Biostratigraphy*. Kluwer Academic Publisher, Dordrecht, Boston, London, pp. 132–199.
- Leonhard, R. (1897) Die Fauna der Kreideformation in Oberschlesien. *Palaeontographica* 44(1-2): 11–70.
- Kear, B.P., Ekrt, B., Prokop J. & Georgalis, G.L. (2014) Turonian marine amniotes from the Bohemian Cretaceous Basin, Czech Republic. *Geological Magazine* 151(1):183–198.
- Madzia, D. (2015) The dentition of *Polyptychodon interruptus* (Plesiosauria, Pliosauridae). In *European Association of Vertebrate Palaeontologists, XIII Annual Meeting, 8–12 July 2015, Abstracts*: 130.
- Sachs, S., Wilmsen, M., Knüppe, J., Hornung, J.J. & Kear, B.P. (2016) Cenomanian–Turonian marine amniote remains from the Saxonian Cretaceous Basin of Germany. *Geological Magazine*. 1021 DOI:10.1017/S0016756815001004.
-

5th Triennial Mosasaur Meeting



A global perspective on Mesozoic marine amniotes


May 16–20, 2016
Museum of Evolution, Uppsala University
Uppsala, Sweden

Program and Abstracts

Benjamin P. Kear, Johan Lindgren & Sven Sachs, Editors



**UPPSALA
UNIVERSITET**



namu

natur | mensch | umwelt